

Dependence on the parameters of the solution of a mixed problem for a nonlinear integro-differential equation of the fifth order with a degenerate kernel

Tursun K. Yuldashev, Khanlar R. Mamedov, Mahkambek M. Babayev

Abstract. In this paper, it is considered a fifth order nonlinear partial integro-differential equations with mixed conditions and two real parameters. The Fourier spectral method of separation of variables is applied. A countable system of nonlinear functional-integral equations is derived. Theorem on a uniqueness and existence of the solution of mixed problem is proved for regular values of parameters. The method of compressing mapping in Banach space is applied. The solution of the mixed problem is obtained in the form of Fourier series. Theorem on absolute and uniform convergence of Fourier series is proved. Continuous dependence on parameters of the classical solution of mixed problem is studied.

Mixed problem, fifth order differential equation, unique solvability, real parameters, regular values, dependence on the parameters.

2010 Mathematics Subject Classifications: Primary 34B08, 34B10; Secondary 35G05.

1. Formulation of the problem statement

Differential equations of parabolic and hyperbolic types are the base of the equations of mathematical physics. Along with equations of parabolic and hyperbolic types, so-called pseudoparabolic and pseudohyperbolic differential equations are often studied.

Let us consider differential equations of the form

$$\left[\frac{\partial^k}{\partial t^k} + (-1)^m \frac{\partial^{k+2m}}{\partial t^k \partial x^{2m}} + (-1)^m \omega \frac{\partial^{2m}}{\partial x^{2m}} \right] U(t, x) = f(t, x), \quad k = 1, 2, 3, \quad m = 1, 2, 3, \dots, n.$$

This equation is sometimes called a Barenblatt–Zheltev–Kochina equation at $k = 1$. And when $k = 2$, it is often called a Boussinesq type differential equation. Many works have been devoted to the study of this equation for $k = 1, 2$ (see, for example, [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51]). However, we have not yet encountered a single work on the study of an equation in the case of $k = 3$.

Integro-differential equations are studied in the works of many mathematicians (see, for examples [52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78]). To study integro-differential equations with degenerate kernel are devoted only the few works [79, 80, 81, 82, 83, 84, 85, 86, 87, 88].

In this paper we consider the case $k = 3$. For simplicity, we take $m = 1$. So, we study the solvability of the mixed problem for a fifth order integro-differential equation with two real parameter and degenerate kernel. So, in the rectangular domain, $\Omega = \{0 < t < T, 0 < x < 1\}$ we consider the following partial integro-differential equation

$$\begin{aligned} & \left[\frac{\partial^3}{\partial t^3} - \frac{\partial^5}{\partial t^3 \partial x^2} - \omega \frac{\partial^2}{\partial x^2} \right] V(t, x) = \\ & = \nu \int_0^T K(t, s) V(s, x) ds + F \left(t, x, \int_0^T \int_0^1 G(s, y) V(s, y) dy ds \right), \end{aligned} \quad (1)$$

where $K(t, s) = \sum_{i=1}^p \alpha_i(t) \beta_i(s)$, $0 < \alpha_i(t)$, $\beta_i(s) \in C[0, T]$, $\alpha_i(t)$ and $\beta_i(s)$ are linear independent, $F(t, x, y) \in C_{t,x,u}^{0,2,0}(\bar{\Omega} \times R)$, $0 < G(t, x) \in C(\bar{\Omega})$, T is given positive number, ω is positive finite parameter, ν is nonzero real parameter, $\bar{\Omega} = \{0 \leq t \leq T, 0 \leq x \leq 1\}$.

In solving partial integro-differential equation (1), we use the following spectral and initial value conditions

$$V(t, 0) = V(t, 1) = 0, \quad 0 \leq t \leq T, \quad (2)$$

$$V(0, x) = \varphi_1(x), \quad V_t(0, x) = \varphi_2(x), \quad V_{tt}(0, x) = \varphi_3(x), \quad 0 \leq x \leq 1, \quad (3)$$

where $\varphi_k(x)$ are enough smooth functions on the segment $[0, 1]$. For these functions the following conditions are fulfilled $\varphi_k(0) = \varphi_k(1) = 0$, $k = 1, 2, 3$.

Problem statement. To find a function

$$V(t, x) \in C(\bar{\Omega}) \cap C_{t,x}^{3,2}(\Omega), \quad (4)$$

which satisfies integro-differential equation (1) and conditions (2) and (3).

2. Countable system of nonlinear equations

The solution of the problem (1)–(3) we search in the form of Fourier series

$$V(t, x) = \sum_{n=1}^{\infty} a_n(t) b_n(x), \quad (5)$$

where $b_n(x) = \sqrt{2} \sin \sqrt{\lambda_n} x$ are eigenfunctions of the spectral problem $b''(x) + \lambda b(x) = 0$, $b(0) = b(1) = 0$, corresponding to the eigenvalues $\lambda_n = (n\pi)^2$,

$$a_n(t) = \int_0^1 V(t, x) b_n(x) dx.$$

So, we assume that

$$F(t, x, y) = \sum_{n=1}^{\infty} F_n(t, \cdot) b_n(x), \quad (6)$$

where

$$F_n(t, \cdot) = \int_0^1 F(t, y, \cdot) b_n(y) dy.$$

Substituting the Fourier series (5) and (6) into the given integro-differential equation (1), and taking into account that the functions $\{b_n(x)\}_{n=1}^{\infty}$ form a complete system of orthonormal systems, we obtain a countable system of third order ordinary differential equations

$$a_n'''(t) + \mu_n(\omega)a_n(t) = f_n(t), \quad \mu_n(\omega) = \frac{\lambda_n}{1 + \lambda_n}\omega, \quad \lambda_n = (n\pi)^2, \quad (7)$$

where

$$f_n(t) = \frac{1}{1 + \lambda_n} \left[\nu \sum_{i=1}^p \alpha_i(t) \tau_{n,i} + F_n(t, \cdot) \right], \quad (8)$$

$$\tau_{n,i} = \int_0^T \beta_i(s) a_n(s) ds. \quad (9)$$

The characteristic equation $\sigma^3 + \mu_n(\omega) = 0$ for the homogeneous equation $a_n'''(t) + \mu_n(\omega)a_n(t) = 0$ has the roots

$$\sigma_1 = -\sqrt[3]{\mu_n(\omega)}, \quad \sigma_{2/3} = \left(\frac{1}{2} \pm \frac{\sqrt{3}}{2}i \right) \sqrt[3]{\mu_n(\omega)}.$$

So, the general solution of the homogeneous equation can be presented as

$$a_n(t) = A_{1,n}a_{1,n}(t) + A_{2,n}a_{2,n}(t) + A_{3,n}a_{3,n}(t), \quad (10)$$

where $A_{k,n}$ ($k = 1, 2, 3$) are yet arbitrary coefficients, which will be determined later,

$$\begin{cases} a_{1,n}(t) = e^{-\sqrt[3]{\mu_n(\omega)}t}, \\ a_{2,n}(t) = e^{\frac{\sqrt[3]{\mu_n(\omega)}t}{2}} \cos \frac{\sqrt{3}}{2} \sqrt[3]{\mu_n(\omega)}t, \\ a_{3,n}(t) = e^{\frac{\sqrt[3]{\mu_n(\omega)}t}{2}} \sin \frac{\sqrt{3}}{2} \sqrt[3]{\mu_n(\omega)}t. \end{cases} \quad (11)$$

Taking (10) into account, we search a particular solution of the equation (7) as

$$\tilde{a}_n(t) = A_{1,n}(t)a_{1,n}(t) + A_{2,n}(t)a_{2,n}(t) + A_{3,n}(t)a_{3,n}(t). \quad (12)$$

In (12) we supposed that $A_{k,n}(t)$ are unknown functions. To find these functions we consider the following system of algebraic-differential equations

$$\begin{cases} A'_{1,n}(t)a_{1,n}(t) + A'_{2,n}(t)a_{2,n}(t) + A'_{3,n}(t)a_{3,n}(t) = 0, \\ A'_{1,n}(t)a'_{1,n}(t) + A'_{2,n}(t)a'_{2,n}(t) + A'_{3,n}(t)a'_{3,n}(t) = 0, \\ A'_{1,n}(t)a''_{1,n}(t) + A'_{2,n}(t)a''_{2,n}(t) + A'_{3,n}(t)a''_{3,n}(t) = f_n(t). \end{cases}$$

We solve the system as functional-algebraic equations by the Cramer rule and found:

$$A_{1,n}(t) = \frac{1}{12c_n^2} \int_0^t \frac{1}{a_{1,n}(s)} f_n(s) ds, \quad (13)$$

$$A_{2,n}(t) = -\frac{1}{12c_n^2} \int_0^t \left[a_{1,n}(s)a_{2,n} + \sqrt{3}a_{1,n}(s)a_{3,n}(s) \right] f_n(s) ds, \quad (14)$$

$$A_{3,n}(t) = \frac{1}{12c_n^2} \int_0^t \left[\sqrt{3}a_{1,n}(s)a_{2,n}(s) - a_{1,n}(s)a_{3,n}(s) \right] f_n(s) ds, \quad (15)$$

where $2c_n = \sqrt[3]{\mu_n(\omega)}$. Substituting (13)–(15) into (12) and taking into account (11), we obtain a particular solution of the equation (7) as

$$\tilde{a}_n(t, \omega) = \frac{1}{\sqrt[3]{\mu_n^2(\omega)}} \int_0^t Q(t, s, \omega) f_n(s) ds, \quad (16)$$

where

$$Q(t, s, \omega) = \frac{1}{3} \left\{ e^{-\sqrt[3]{\mu_n(\omega)}(t-s)} - 2e^{\frac{\sqrt[3]{\mu_n(\omega)}}{2}(t-s)} \sin \left(\frac{\sqrt{3}}{2} \sqrt[3]{\mu_n(\omega)}(s-t) + \frac{\pi}{6} \right) \right\}.$$

From (16) we derive a particular solution of the equation (1) in the form of Fourier series

$$\tilde{V}(t, x, \omega) = \sum_{n=1}^{\infty} \frac{b_n(x)}{\sqrt[3]{\mu_n^2(\omega)}} \int_0^t Q(t, s, \omega) f_n(s) ds. \quad (17)$$

Taking into account (16) the general solution of the equation (7) can be presented as:

$$\begin{aligned} a_n(t) &= B_{1,n}a_{1,n}(t) + B_{2,n}a_{2,n}(t) + B_{3,n}a_{3,n}(t) + \\ &+ A_{1,n}(t)a_{1,n}(t) + A_{2,n}(t)a_{2,n}(t) + A_{3,n}(t)a_{3,n}(t) \end{aligned} \quad (18)$$

and we find the constants $B_{k,n}$ ($k = 1, 2, 3$).

We suppose that the functions $\varphi_k(x)$ are expanding into a Fourier series and using the Fourier coefficients, from initial value conditions (3) we obtain

$$a_n(0) = \int_0^1 V(0, y) b_n(y) dy = \int_0^1 \varphi_1(y) b_n(y) dy = \varphi_{1,n}, \quad (19)$$

$$a'_n(0) = \int_0^1 V_t(0, y) b_n(y) dy = \int_0^1 \varphi_2(y) b_n(y) dy = \varphi_{2,n}, \quad (20)$$

$$a''_n(0) = \int_0^1 V_{tt}(0, y) b_n(y) dy = \int_0^1 \varphi_3(y) b_n(y) dy = \varphi_{3,n}. \quad (21)$$

To find the unknown (arbitrary) coefficients $A_{k,n}$ ($k = 1, 2, 3$), we use the boundary conditions (19)–(21) in the presentation (18) and obtain

$$\begin{cases} B_{1,n}a_{1,n} = \frac{1}{3}a_{1,n}\varphi_{1,n} - \frac{1}{6b_n}a_{1,n}\varphi_{2,n} + \frac{1}{12b_n^2}a_{1,n}\varphi_{3,n}, \\ B_{2,n}a_{2,n} = \frac{2}{3}a_{2,n}\varphi_{1,n} + \frac{1}{6b_n}a_{2,n}\varphi_{2,n} - \frac{1}{12b_n^2}a_{2,n}\varphi_{3,n}, \\ B_{3,n}a_{3,n} = \frac{1}{2\sqrt{3}c_n}a_{3,n}\varphi_{2,n} + \frac{1}{4\sqrt{3}c_n^2}a_{3,n}\varphi_{3,n}. \end{cases}$$

Hence, we obtain that

$$\begin{aligned} & B_{1,n}a_{1,n} + B_{2,n}a_{2,n} + B_{3,n}a_{3,n} = \\ & = \frac{a_{1,n} + 2a_{2,n}}{3}\varphi_{1,n} + \frac{-a_{1,n} + a_{2,n} + \sqrt{3}a_{3,n}}{6c_n}\varphi_{2,n} + \frac{a_{1,n} - c_{2,n} + \sqrt{3}c_{3,n}}{12c_n^2}\varphi_{3,n}. \end{aligned} \quad (22)$$

Substituting (22) into (18) and taking into account (8), (11), (16), we obtain

$$\begin{aligned} a_n(t, \omega) &= P_n(t, \omega) + \\ &+ \frac{1}{\sqrt[3]{(\lambda_n^2 + \lambda_n^3)\omega}} \left[\nu \sum_{i=1}^p \tau_{n,i} \int_0^t Q_n(t, s, \omega) \alpha_i(s) ds + \int_0^t Q_n(t, s, \omega) F_n(s, \cdot) ds \right], \end{aligned} \quad (23)$$

where

$$P_n(t, \omega) = \varphi_{1,n}\psi_{1,n}(t, \omega) + \frac{1}{\sqrt[3]{\mu_n(\omega)}}\varphi_{2,n}\psi_{2,n}(t, \omega) + \frac{1}{\sqrt[3]{\mu_n^2(\omega)}}\varphi_{3,n}\psi_{3,n}(t, \omega), \quad (24)$$

$$Q_n(t, s, \omega) = \frac{1}{3} \left[e^{-\sqrt[3]{\mu_n(\omega)}(t-s)} + 2e^{\frac{\sqrt[3]{\mu_n(\omega)}}{2}(t-s)} \sin \left(\frac{\sqrt{3}}{2} \sqrt[3]{\mu_n(\omega)}(t-s) + \frac{\pi}{6} \right) \right], \quad (25)$$

$$\psi_{1,n}(t, \omega) = \frac{1}{3} \left[e^{-\sqrt[3]{\mu_n(\omega)}t} + 2e^{\frac{\sqrt[3]{\mu_n(\omega)}}{2}t} \cos \frac{\sqrt{3}}{2} \sqrt[3]{\mu_n(\omega)}t \right], \quad (26)$$

$$\psi_{2,n}(t, \omega) = \frac{1}{3} \left[e^{-\sqrt[3]{\mu_n(\omega)}t} - 2e^{\frac{\sqrt[3]{\mu_n(\omega)}}{2}t} \sin \left(\frac{\sqrt{3}}{2} \sqrt[3]{\mu_n(\omega)}t + \frac{\pi}{6} \right) \right], \quad (27)$$

$$\psi_{3,n}(t, \omega) = \frac{1}{3} \left[e^{-\sqrt[3]{\mu_n(\omega)}t} - 2e^{\frac{\sqrt[3]{\mu_n(\omega)}}{2}t} \sin \left(\frac{\sqrt{3}}{2} \sqrt[3]{\mu_n(\omega)}t - \frac{\pi}{6} \right) \right], \quad (28)$$

$$F_n(t, \cdot) = \int_0^1 F \left(t, y, \int_0^T \int_0^1 G(s, z) V(s, z) dz ds \right) b_n(y) dy.$$

There is another unknown quantity in (23). To find it we substitute (23) into (9), and obtain a countable system of algebraic system of equations (CSASE)

$$\tau_{n,i} = \nu \sum_{j=1}^p \tau_{n,j} \Phi_{n,i,j}(\omega) + \Psi_{n,i}(a_r, \omega), \quad i = 1, 2, \dots, p, \quad (29)$$

$$\Phi_{n,i,j}(\omega) = \frac{1}{\sqrt[3]{(\lambda_n^2 + \lambda_n^3)\omega}} \int_0^T \beta_i(s) \int_0^s Q_n(s, \theta, \omega) \alpha_j(\theta) d\theta ds, \quad (30)$$

$$\begin{aligned} \Psi_{n,i}(a_r, \omega) &= \int_0^T \beta_i(s) P_n(s, \omega) ds + \frac{1}{\sqrt[3]{(\lambda_n^2 + \lambda_n^3)\omega}} \int_0^T \beta_i(s) \int_0^s Q_n(s, \theta, \omega) \times \\ &\times \int_0^1 F \left(\theta, y, \int_0^T \int_0^1 G(\xi, z) \sum_{r=1}^{\infty} a_r(\xi) b_r(z) dz d\xi \right) b_n(y) dy d\theta ds. \end{aligned} \quad (31)$$

To solve the CSASE we consider the following determinants

$$Z_n(\nu, \omega) = \begin{vmatrix} 1 - \nu \Phi_{11} & \nu \Phi_{12} & \dots & \nu \Phi_{1p} \\ \nu \Phi_{21} & 1 - \nu \Phi_{22} & \dots & \nu \Phi_{2p} \\ \dots & \dots & \dots & \dots \\ \nu \Phi_{p1} & \nu \Phi_{p2} & \dots & 1 - \nu \Phi_{pp} \end{vmatrix}, \quad (32)$$

$$Z_{in}(a_r, \nu, \omega) = \begin{vmatrix} 1 - \nu \Phi_{11} & \dots & \nu \Phi_{1(i-1)} & \Psi_1 & \nu \Phi_{1(i+1)} & \dots & \nu \Phi_{1p} \\ \nu \Phi_{21} & \dots & \nu \Phi_{2(i-1)} & \Psi_2 & \nu \Phi_{2(i+1)} & \dots & \nu \Phi_{2p} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \nu \Phi_{p1} & \dots & \nu \Phi_{p(i-1)} & \Psi_p & \nu \Phi_{p(i+1)} & \dots & 1 - \nu \Phi_{pp} \end{vmatrix}, \quad (33)$$

where $\Phi_{ij} = \Phi_{nij}(\omega)$, $\Psi_\kappa = \Psi_{n\kappa}(a_r, \omega)$, $\kappa = \overline{1, p}$.

CSASE (29) is uniquely solvable for any finite right-hand sides, if the following non-degeneracy condition for the Fredholm determinant is satisfied $Z_n(\nu, \omega) \neq 0$. The determinant (32) $Z_n(\nu, \omega)$ is a polynomial with respect to ν of degree not higher p . The equation $Z_n(\nu, \omega) = 0$ has at most p different real roots. We denote them by θ_ℓ ($\ell = \overline{1, p}$, $1 \leq p_\ell \leq p$). Then $\nu = \nu_{n+\ell} = \theta_\ell$ called the irregular values of the parameter

ν . Other values of the parameter $\nu \neq \theta_\ell$, for which $|Z_n(\nu, \omega)| > 0$ and $\sum_{n=1}^{\infty} \frac{1}{[Z_n(\nu, \omega)]^2} < \infty$, are called regular.

For regular values of the parameter ν the solution of the CSASE (29) has the form

$$\tau_{n\kappa}(\nu, \omega) = \frac{Z_{\kappa n}(a_r, \nu, \omega)}{Z_n(\nu, \omega)}, \quad \kappa = \overline{1, p}, \quad (34)$$

where $Z_{\kappa n}(\nu, \omega)$ is defined from (33). Substituting for regular values of the parameter ν the presentation of solution (34) of the CSASE (29) into representation (23) of the Fourier coefficients $a_n(t)$ of unknown function $V(t, x, \nu, \omega)$, we derive a nonlinear countable system of functional integral equations (NCSFIE)

$$\begin{aligned} a_n(t, \nu, \omega) &= J(t; a_n) \equiv P_n(t, \nu, \omega) + \\ &+ \frac{\nu}{\sqrt[3]{(\lambda_n^2 + \lambda_n^3)\omega}} \sum_{\kappa=1}^p \frac{Z_{\kappa n}(a_r, \nu, \omega)}{Z_n(\nu, \omega)} \int_0^t Q_n(t, s, \omega) \alpha_j(s) ds + \frac{1}{\sqrt[3]{(\lambda_n^2 + \lambda_n^3)\omega}} \times \\ &\times \int_0^t Q_n(t, s, \omega) \int_0^1 F \left(s, y, \int_0^T \int_0^1 G(\theta, z) \sum_{r=1}^{\infty} a_r(\theta) b_r(z) dz d\theta \right) b_n(y) dy ds. \end{aligned} \quad (35)$$

As, in the case of Fourier series (17), from (35) we obtain a formal solution of the mixed problem (1)–(3)

$$\begin{aligned} V(t, x, \nu, \omega) &= \sum_{n=1}^{\infty} b_n(x) \left[P_n(t, \omega) + \right. \\ &+ \frac{\nu}{\sqrt[3]{(\lambda_n^2 + \lambda_n^3)\omega}} \sum_{\kappa=1}^p \frac{Z_{\kappa n}(a_r, \nu, \omega)}{Z_n(\nu, \omega)} \int_0^t Q_n(t, s, \omega) \alpha_j(s) ds + \frac{1}{\sqrt[3]{(\lambda_n^2 + \lambda_n^3)\omega}} \times \\ &\times \left. \int_0^t Q_n(t, s, \omega) \int_0^1 F \left(s, y, \int_0^T \int_0^1 G(\theta, z) \sum_{r=1}^{\infty} a_r(\theta) b_r(z) dz d\theta \right) b_n(y) dy ds \right]. \end{aligned} \quad (36)$$

We note that the functions in (25)–(28) become zero at some values of parameter ω . We obtain the following transcendental equation

$$\sin \left(\frac{\sqrt{3}}{2} y + \frac{\pi}{6} \right) = -\frac{1}{2} e^{-\frac{3}{2} y}, \quad y = \sqrt[3]{\mu_n(\omega)} (t - s) > 0,$$

for the case of function (25) and

$$\cos \frac{\sqrt{3}}{2} y = -\frac{1}{2} e^{-\frac{3}{2} y}, \quad y = \sqrt[3]{\mu_n(\omega)} t > 0,$$

for the case of function (26), respectively, $\mu_n(\omega) = \frac{\lambda_n}{1+\lambda_n}\omega$. Functions in the formulas (27) and (28) become zero at some values of parameter ω . We replace these equations by the following transcendental equations

$$\sin\left(\frac{\sqrt{3}}{2}y + \frac{\pi}{6}\right) = \frac{1}{2}e^{\frac{-3y}{2}}, \quad y = \sqrt[3]{\mu_n(\omega)} \quad t > 0,$$

$$\sin\left(\frac{\sqrt{3}}{2}y - \frac{\pi}{6}\right) = \frac{1}{2}e^{\frac{-3y}{2}}, \quad y = \sqrt[3]{\mu_n(\omega)} \quad t > 0,$$

respectively.

The values of parameter ω , for which the functions (25)–(28) become zero, we denote by Λ_j , $j = 1, 2, 3, 4$, respectively. However, from the fact $\Lambda_1 \cap \Lambda_2 \cap \Lambda_3 \cap \Lambda_4 = \emptyset$ we deduce that the problem (1)–(3) is correct.

3. Solvability of the countable system

To prove the classical solvability of the mixed problem (1)–(3) we require in some properties of the given functions, which we call as smoothness conditions.

Smoothness conditions. Let the functions $\varphi_k(x) \in C^4[0, 1]$ ($k = 1, 2, 3$), $F(t, x, \cdot) \in C_{t,x}^{0,2}(\Omega \times R)$ have continuous derivatives with respect to x up to the fourth and second order, respectively. We integrate by parts

$$\varphi_{k,n} = \int_0^1 \varphi_k(y) b_n(y) dy, \quad F_n(t, \cdot) = \int_0^1 F\left(t, y, \int_0^T \int_0^1 G(s, z) V(s, z) dz ds\right) b_n(y) dy$$

four and second times, respectively. Then we obtain the estimates

$$\varphi_{k,n} \leq \left(\frac{1}{\pi}\right)^4 \frac{|\varphi_{k,n}^{(IV)}|}{n^4}, \quad F_n \leq \left(\frac{1}{\pi}\right)^2 \frac{|F_n''(t, \cdot)|}{n^2},$$

where

$$\varphi_{k,n}^{(IV)} = \int_0^1 \frac{\partial^4 \varphi_k(y)}{\partial y^4} b_n(y) dy, \quad F_n''(t, \cdot) = \int_0^1 \frac{\partial^2}{\partial y^2} F\left(t, y, \int_0^T \int_0^1 G(s, z) V(s, z) dz ds\right) b_n(y) dy.$$

In estimating approximations, we use also Bessel inequalities in the form:

$$\left\| \varphi_k^{(IV)} \right\|_{\ell_2} \leq 4 \left\| \frac{\partial^4 \varphi_k(x)}{\partial x^4} \right\|_{L_2[0,1]}, \quad \left\| \vec{F}''(t, \cdot) \right\|_{B_2[0,T]} \leq 2 \max_{0 \leq t \leq T} \left\| \frac{\partial^4 F(t, x, \cdot)}{\partial x^4} \right\|_{L_2[0,1]}.$$

Theorem 3.1. *Let the smoothness conditions be fulfilled and*

- 1). $\alpha_0 \sum_{\kappa=1}^p |Z_{\kappa n}(a_r^0, \nu, \omega)| \leq \delta_0$, $\alpha_0 = \max_{0 \leq t \leq T} \int_0^t \alpha_j(s) ds$, $0 < \delta_0$, $\alpha_0 = \text{const} < \infty$;

- 2). $\max_{0 \leq t \leq T} \|F(t, x, \cdot)\|_{L_2[0,1]} \leq \delta_1$, $0 < \delta_1 = \text{const} < \infty$;
3). $|F(t, x, u_1) - F(t, x, u_2)| \leq l(x) |u_1 - u_2|$, $0 < l(x) \in L_2[0, 1]$;
4). $\rho = M_3 \int_0^T \|G(t, x)\|_{L_2[0,1]} dt < 1$, where M_3 determines from (41) below.

Then for regular values of the parameter ν NCSFIE (35) has a unique solution in the space $B_2[0, T]$ with norm

$$\|\vec{a}(t)\|_{B_2[0,T]} = \sqrt{\sum_{n=1}^{\infty} \left(\max_{t \in [0, T]} |a_n(t)| \right)^2} < \infty.$$

Proof. We define the successive approximations for NCSFIE (35) as:

$$\begin{cases} a_n^0(t, \nu, \omega) = P_n(t, \nu, \omega), \\ a_n^{m+1}(t, \nu, \omega) = J(t; a_n^m), \quad m = 1, 2, 3, \dots \end{cases} \quad (37)$$

We estimate the zero approximation. By virtue of formulas (24), (26)–(28), and

$$\mu_n = \frac{\lambda_n}{1 + \lambda_n} < 1, \quad \lim_{\lambda_n \rightarrow \infty} \frac{1}{\sqrt[3]{\mu_n^2(\omega)}} = \lim_{\lambda_n \rightarrow \infty} \sqrt[3]{\left(\frac{1 + \lambda_n}{\lambda_n}\right)^2 \frac{1}{\omega^2}} = e^{\frac{2}{3}} \omega^{-\frac{2}{3}}, \quad \omega > 0,$$

we can put

$$\max \left\{ \max_t |Q_n(t, s, \omega)|; \max_{k=1,2,3} \max_t |\psi_{k,n}(t, \omega)| \right\} \leq M_0 < \infty, \quad 0 < M_0 = \text{const} < \infty.$$

By virtue of smoothness conditions, applying the Cauchy–Schwartz inequality and Bessel inequality, from approximations (37) we have

$$\begin{aligned} \|\vec{a}^0(t, \nu, \omega)\|_{B_2[0,T]} &\leq \sqrt{\sum_{n=1}^{\infty} \left[\max_{0 \leq t \leq T} |a_n^0(t, \nu, \omega)| \right]^2} \leq \\ &\leq \sum_{n=1}^{\infty} \max_{0 \leq t \leq T} |P_n(t, \omega)| \leq M_0 \left[\sum_{n=1}^{\infty} |\varphi_{1,n}| + \sqrt[3]{\frac{e}{\omega}} \sum_{n=1}^{\infty} |\varphi_{2,n}| + \sqrt[3]{\left(\frac{e}{\omega}\right)^2} \sum_{n=1}^{\infty} |\varphi_{3,n}| \right] \leq \\ &\leq C_0 M_0 \left(\frac{1}{\pi}\right)^4 \sqrt{\sum_{n=1}^{\infty} \frac{1}{n^8} \left[\|\vec{\varphi}_1^{(IV)}\|_{\ell_2} + \|\vec{\varphi}_2^{(IV)}\|_{\ell_2} + \|\vec{\varphi}_3^{(IV)}\|_{\ell_2} \right]} \leq \\ &\leq M_1 \left[\left\| \frac{\partial^4 \varphi_1(x)}{\partial x^4} \right\|_{L_2[0,1]} + \left\| \frac{\partial^4 \varphi_2(x)}{\partial x^4} \right\|_{L_2[0,1]} + \left\| \frac{\partial^4 \varphi_3(x)}{\partial x^4} \right\|_{L_2[0,1]} \right] < \infty, \quad (38) \end{aligned}$$

where

$$M_1 = C_0 M_0 \left(\frac{\sqrt{2}}{\pi}\right)^4 \sqrt{\sum_{n=1}^{\infty} \frac{1}{n^8}}, \quad C_0 = \max \left\{ 1; \sqrt[3]{\frac{e}{\omega}}; \sqrt[3]{\left(\frac{e}{\omega}\right)^2} \right\}.$$

Due to the conditions of the Theorem 3.1, formulas (25), (30), (31), estimate (38) and applying the Cauchy–Schwartz inequality and Bessel inequality, for the first difference $a_n^1(t) - a_n^0(t)$ we obtain

$$\begin{aligned}
& \left\| \bar{a}^1(t, \nu, \omega) - \bar{a}^0(t, \nu, \omega) \right\|_{B_2[0, T]} \leq \\
& \leq \frac{|\nu|}{\sqrt[3]{\omega}} \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \sum_{\kappa=1}^p \left| \frac{Z_{\kappa n}(a_r^0, \nu, \omega)}{Z_n(\nu, \omega)} \right| \max_{0 \leq t \leq T} \int_0^t |Q_n(t, s, \omega)| \alpha_j(s) ds + \\
& + \frac{1}{\sqrt[3]{\omega}} \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \max_{0 \leq t \leq T} \int_0^t |Q_n(t, s, \omega)| \left| \int_0^1 F \left(s, y, \int_0^T \int_0^1 G(\theta, z) \sum_{r=1}^{\infty} a_r^0(\theta, \omega) b_r(z) dz d\theta \right) b_n(y) dy \right| ds \leq \\
& \leq \frac{M_0}{\sqrt[3]{\omega}} \left(\frac{1}{\pi} \right)^2 \sqrt{\sum_{n=1}^{\infty} \frac{1}{n^4}} \left[|\nu| \delta_0 \sqrt{\sum_{n=1}^{\infty} \frac{1}{|Z_n(\nu, \omega)|^2}} + T \left\| \int_0^1 F(t, y, \cdot) b_n(y) dy \right\|_{B_2[0, T]} \right] \leq \\
& \leq M_2 \left[|\nu| \delta_0 \sqrt{\sum_{n=1}^{\infty} \frac{1}{|Z_n(\nu, \omega)|^2}} + T \max_{0 \leq t \leq T} \|F(t, x, \cdot)\|_{L_2[0, 1]} \right] < \infty, \quad (39)
\end{aligned}$$

where $M_2 = \frac{M_0}{\sqrt[3]{\omega}} \left(\frac{1}{\pi} \right)^2 \sqrt{\sum_{n=1}^{\infty} \frac{1}{n^4}}$.

Now we consider the arbitrary consecutive difference $a_n^{m+1}(t) - a_n^m(t)$. We take into account that the quantities

$$\int_0^1 l(y) b_n(y) dy, \quad \int_0^1 |G(t, z)| b_r(z) dz$$

are Fourier coefficients. By the same way as the estimate (39) above, we obtain

$$\begin{aligned}
& \left\| \bar{a}^{m+1}(t, \nu, \omega) - \bar{a}^m(t, \nu, \omega) \right\|_{B_2[0, T]} \leq \\
& \leq \frac{|\nu|}{\sqrt[3]{\omega}} \sum_{n=1}^{\infty} \frac{M_0}{\lambda_n} \sum_{i=1}^p \left| \frac{Z_{in}(a_r^m, \nu, \omega) - Z_{in}(a_r^{m-1}, \nu, \omega)}{Z_n(\nu, \omega)} \right| \max_{0 \leq t \leq T} \int_0^t \alpha_j(s) ds + \\
& + M_0 \frac{1}{\sqrt[3]{\omega}} \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \max_{0 \leq t \leq T} \int_0^t \left| \int_0^1 l(y) \int_0^T \int_0^1 |G(t, z)| \sum_{r=1}^{\infty} |a_r^m(t, \omega) - a_r^{m-1}(t, \omega)| b_r(z) dz dt b_n(y) dy \right| ds \leq \\
& \leq \frac{|\nu|}{\sqrt[3]{\omega^2}} \sum_{n=1}^{\infty} \frac{\alpha_0 M_0^2}{\lambda_n^2 Z_n(\nu, \omega)} \sum_{\kappa=1}^p \bar{\delta}_{0, \kappa} \int_0^T \beta_{\kappa}(s) ds
\end{aligned}$$

$$\begin{aligned}
& \left| \int_0^1 l(y) b_n(y) dy \int_0^T \int_0^1 |G(t, z)| \sum_{r=1}^{\infty} |a_r^m(t, \omega) - a_r^{m-1}(t, \omega)| b_r(z) dz dt \right| + \\
& + M_0 \frac{T}{\sqrt[3]{\omega}} \left| \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \int_0^1 l(y) b_n(y) dy \int_0^T \int_0^1 |G(t, z)| \sum_{r=1}^{\infty} |a_r^m(t, \omega) - a_r^{m-1}(t, \omega)| b_r(z) dz dt \right| \leq \\
& \leq \frac{|\nu|}{\sqrt[3]{\omega^2}} \sum_{n=1}^{\infty} \frac{\alpha_0 \beta_0 M_0^2}{\lambda_n^2 Z_n(\nu, \omega)} \left| \int_0^1 l(y) b_n(y) dy \right| \left| \int_0^T \int_0^1 |G(t, z)| \sum_{r=1}^{\infty} |a_r^m(t, \omega) - a_r^{m-1}(t, \omega)| b_r(z) dz dt \right| + \\
& + M_0 \frac{T}{\sqrt[3]{\omega}} \left| \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \int_0^1 l(y) b_n(y) dy \right| \left| \int_0^T \int_0^1 |G(t, z)| \sum_{r=1}^{\infty} |a_r^m(t, \omega) - a_r^{m-1}(t, \omega)| b_r(z) dz dt \right| \leq \\
& \leq \alpha_0 \beta_0 \delta_2 M_0^2 \frac{|\nu|}{\sqrt[3]{\omega^2}} \left\| \frac{1}{\bar{\lambda}^4 \bar{Z}^2(\nu, \omega)} \right\|_{\ell_2} \left| \int_0^T \int_0^1 |G(t, z)| \sum_{r=1}^{\infty} |a_r^m(t, \omega) - a_r^{m-1}(t, \omega)| b_r(z) dz dt \right| + \\
& + M_0 \delta_2 \frac{T}{\sqrt[3]{\omega}} \left\| \frac{1}{\bar{\lambda}^2} \right\|_{\ell_2} \left| \int_0^T \int_0^1 |G(t, z)| \sum_{r=1}^{\infty} |a_r^m(t, \omega) - a_r^{m-1}(t, \omega)| b_r(z) dz dt \right| \leq \\
& \leq M_3 \int_0^T \sum_{r=1}^{\infty} \left| \int_0^1 |G(t, z)| b_r(z) dz \right| |a_r^m(t, \nu, \omega) - a_r^{m-1}(t, \nu, \omega)| dt \leq \\
& \leq \rho \cdot \left\| \bar{a}^m(t, \nu, \omega) - \bar{a}^{m-1}(t, \nu, \omega) \right\|_{B_2[0, T]}, \tag{40}
\end{aligned}$$

where

$$\rho = M_3 \int_0^T \|G(t, x)\|_{L_2[0, 1]} dt, \tag{41}$$

$$M_3 = \alpha_0 \beta_0 \delta_2 M_0^2 \frac{|\nu|}{\sqrt[3]{\omega^2}} \left\| \frac{1}{\bar{\lambda}^4 \bar{Z}^2(\nu, \omega)} \right\|_{\ell_2} + M_0 \delta_2 \frac{T}{\sqrt[3]{\omega}} \left\| \frac{1}{\bar{\lambda}^2} \right\|_{\ell_2},$$

$$\beta_0 = \sum_{\kappa=1}^p \bar{\delta}_{0, \kappa} \int_0^T \beta_{\kappa}(s) ds, \quad \bar{\delta}_{0, \kappa} = |\bar{Z}_{\kappa n}(\nu, \omega)|,$$

$$\delta_2 = \|l(x)\|_{L_2[0, 1]} \geq \sqrt{\sum_{n=1}^{\infty} \left| \int_0^1 l(y) b_n(y) dy \right|^2},$$

$$\bar{Z}_{in}(\nu, \omega) = \begin{vmatrix} 1 - \nu \Phi_{11} & \dots & \nu \Phi_{1(i-1)} & 1 & \nu \Phi_{1(i+1)} & \dots & \nu \Phi_{1p} \\ \nu \Phi_{21} & \dots & \nu \Phi_{2(i-1)} & 1 & \nu \Phi_{2(i+1)} & \dots & \nu \Phi_{2p} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \nu \Phi_{p1} & \dots & \nu \Phi_{p(i-1)} & 1 & \nu \Phi_{p(i+1)} & \dots & 1 - \nu \Phi_{pp} \end{vmatrix}.$$

From estimates (38)–(40) it follows that the operator $J(t; a_n)$ on the right-hand side of (35) is contracting and there is unique fixed point. So, the existence and uniqueness of the solution $\bar{a}(t) \in B_2[0, T]$ to NCSFIE (35) are proved. The theorem 3.1 is proved. \square

4. Continuously dependence of the solution to NCSFIE from parameter ω

In this section we use the following obvious lemma.

Lemma 4.1. *For two values ω_1, ω_2 of positive parameter ω there true the following estimates*

$$\begin{aligned} \left| e^{-\sqrt[3]{\mu_n(\omega_1)}(t-s)} - e^{-\sqrt[3]{\mu_n(\omega_2)}(t-s)} \right| &\leq L_{01} |\omega_1 - \omega_2|, \quad 0 < L_{01} = \text{const}; \\ \left| \sin \left(\frac{\sqrt{3}}{2} \sqrt[3]{\mu_n(\omega_1)}(t-s) + \frac{\pi}{6} \right) - \sin \left(\frac{\sqrt{3}}{2} \sqrt[3]{\mu_n(\omega_2)}(t-s) + \frac{\pi}{6} \right) \right| &\leq L_{02} |\omega_1 - \omega_2|; \\ \left| e^{\sqrt[3]{\mu_n(\omega_1)} \frac{t-s}{2}} - e^{\sqrt[3]{\mu_n(\omega_2)} \frac{t-s}{2}} \right| &\leq L_{03} |\omega_1 - \omega_2|, \quad 0 < L_{03} = \text{const}; \\ \left| \cos \frac{\sqrt{3}}{2} \sqrt[3]{\mu_n(\omega_1)}(t-s) - \cos \frac{\sqrt{3}}{2} \sqrt[3]{\mu_n(\omega_2)}(t-s) \right| &\leq L_{02} |\omega_1 - \omega_2|, \quad 0 < L_{02} = \text{const}; \\ \left| \frac{1}{\sqrt[3]{\omega_1}} - \frac{1}{\sqrt[3]{\omega_2}} \right| &\leq L_{04} |\omega_1 - \omega_2|, \quad L_{04} = \text{const}; \\ \left| \frac{1}{\sqrt[3]{\omega_1^2}} - \frac{1}{\sqrt[3]{\omega_2^2}} \right| &\leq L_{05} |\omega_1 - \omega_2|, \quad L_{05} = \text{const}. \end{aligned}$$

Theorem 4.2. *Let be fulfilled the conditions of the Theorem 3.1. Then the following estimate*

$$\|\bar{a}(t, \omega_1) - \bar{a}(t, \omega_2)\|_{B_2[0, T]} \leq L_M |\omega_1 - \omega_2|, \quad 0 < L_M = \text{const} \quad (42)$$

holds.

Proof. By virtue of the Lemma for the function (25) we obtain

$$\begin{aligned} |Q_n(t, s, \omega_1) - Q_n(t, s, \omega_2)| &\leq \frac{1}{3} \left| e^{-\sqrt[3]{\mu_n(\omega_1)}(t-s)} - e^{-\sqrt[3]{\mu_n(\omega_2)}(t-s)} \right| + \\ + \frac{2}{3} \left| e^{\frac{\sqrt[3]{\mu_n(\omega_1)}}{2}(t-s)} - e^{\frac{\sqrt[3]{\mu_n(\omega_2)}}{2}(t-s)} \right| &\cdot \left| \sin \left(\frac{\sqrt{3}}{2} \sqrt[3]{\mu_n(\omega_1)}(t-s) + \frac{\pi}{6} \right) \right| + \end{aligned}$$

$$\begin{aligned}
& + \frac{2}{3} e^{\frac{\sqrt[3]{\mu_n(\omega_2)}}{2}(t-s)} \left| \sin \left(\frac{\sqrt{3}}{2} \sqrt[3]{\mu_n(\omega_1)}(t-s) + \frac{\pi}{6} \right) - \sin \left(\frac{\sqrt{3}}{2} \sqrt[3]{\mu_n(\omega_2)}(t-s) + \frac{\pi}{6} \right) \right| \leq \\
& \leq \left[L_{01} + L_{03} + e^{\sqrt[3]{\omega_2} T} L_{02} \right] |\omega_1 - \omega_2| = L_1 |\omega_1 - \omega_2|. \tag{43}
\end{aligned}$$

By virtue of the Lemma, for the function (26) we obtain

$$\begin{aligned}
& |\psi_{1,n}(t, \omega_1) - \psi_{1,n}(t, \omega_2)| \leq \\
& \leq \frac{1}{3} \left| e^{-\sqrt[3]{\mu_n(\omega_1)} t} - e^{-\sqrt[3]{\mu_n(\omega_2)} t} \right| + \frac{2}{3} \left| e^{\frac{\sqrt[3]{\mu_n(\omega_1)}}{2} t} - e^{\frac{\sqrt[3]{\mu_n(\omega_2)}}{2} t} \right| \cdot \left| \cos \frac{\sqrt{3}}{2} \sqrt[3]{\mu_n(\omega_1)} t \right| + \\
& + \frac{2}{3} e^{\frac{\sqrt[3]{\mu_n(\omega_2)}}{2} t} \left| \cos \frac{\sqrt{3}}{2} \sqrt[3]{\mu_n(\omega_1)} t - \cos \frac{\sqrt{3}}{2} \sqrt[3]{\mu_n(\omega_2)} t \right| \leq \\
& \leq \left[L_{01} + L_{03} + e^{\sqrt[3]{\omega_2} T} L_{02} \right] |\omega_1 - \omega_2| = L_1 |\omega_1 - \omega_2|. \tag{44}
\end{aligned}$$

By similarly way for the functions (27) and (28) we obtain

$$|\psi_{j,n}(t, \omega_1) - \psi_{j,n}(t, \omega_2)| \leq L_1 |\omega_1 - \omega_2|, \quad j = 2, 3. \tag{45}$$

By the aid of the estimates (43)–(45), taking properties of the functions (24), (31) and matrix (33), for the NCSFIE (35) we derive

$$\begin{aligned}
& \|\vec{a}(t, \nu, \omega_1) - \vec{a}(t, \nu, \omega_2)\|_{B_2[0, T]} \leq \sum_{n=1}^{\infty} \max_{0 \leq t \leq T} |P_n(t, \omega_1) - P_n(t, \omega_2)| + \\
& + |\nu| \left| \frac{1}{\sqrt[3]{\omega_1}} - \frac{1}{\sqrt[3]{\omega_2}} \right| \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \sum_{\kappa=1}^p \left| \frac{Z_{\kappa n}(a_r, \nu, \omega_1)}{Z_n(\nu, \omega_1)} \right| \max_{0 \leq t \leq T} \int_0^t |Q_n(t, s, \omega_1)| \alpha_j(s) ds + \\
& + \frac{|\nu|}{\sqrt[3]{\omega_2}} \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \sum_{\kappa=1}^p \left| \frac{Z_{\kappa n}(a_r, \nu, \omega_1)}{Z_n(\nu, \omega_1)} \right| \max_{0 \leq t \leq T} \int_0^t |Q_n(t, s, \omega_1) - Q_n(t, s, \omega_2)| \alpha_j(s) ds + \\
& + \frac{|\nu|}{\sqrt[3]{\omega_2}} \sum_{n=1}^{\infty} \frac{\alpha_0 M_0}{\lambda_n} \sum_{\kappa=1}^p \left[|Z_{\kappa n}(a_r, \nu, \omega_1)| \left| \frac{1}{Z_n(\nu, \omega_1)} - \frac{1}{Z_n(\nu, \omega_2)} \right| + \right. \\
& \quad \left. + \left| \frac{1}{Z_n(\nu, \omega_2)} \right| |Z_{\kappa n}(a_r, \nu, \omega_1) - Z_{\kappa n}(a_r, \nu, \omega_2)| \right] + \\
& + \left| \frac{1}{\sqrt[3]{\omega_1}} - \frac{1}{\sqrt[3]{\omega_2}} \right| \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \max_{0 \leq t \leq T} \int_0^t Q_n(t, s, \omega_1) \left| \int_0^1 F(s, y, \cdot) b_n(y) dy \right| ds +
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{\sqrt[3]{\omega_2}} \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \max_{0 \leq t \leq T} \int_0^t |Q_n(t, s, \omega_1) - Q_n(t, s, \omega_2)| \left| \int_0^1 F(s, y, \cdot) b_n(y) dy \right| ds + \frac{M_0}{\sqrt[3]{\omega_2}} \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \times \\
& \quad \times \sum_{\kappa=1}^p \left| \int_0^1 l(y) b_n(y) dy \int_0^T \int_0^1 |G(t, z)| \sum_{r=1}^{\infty} |a_r(t, \omega_1) - a_r(t, \omega_2)| b_r(z) dz dt \right|. \quad (46)
\end{aligned}$$

By the aid of Lemma, estimates (43)–(45) and conditions of the Theorem 3.1, from (46) we obtain

$$\begin{aligned}
& \|\bar{a}(t, \nu, \omega_1) - \bar{a}(t, \nu, \omega_2)\|_{B_2[0, T]} \leq C_0 M_0 \left(\frac{\sqrt{2}}{\pi} \right)^4 \sqrt{\sum_{n=1}^{\infty} \frac{1}{n^8}} \left[L_1 \left\| \frac{\partial^4 \varphi_1(x)}{\partial x^4} \right\|_{L_2[0, 1]} + \right. \\
& \quad \left. + (L_1 + L_{04}) \left\| \frac{\partial^4 \varphi_2(x)}{\partial x^4} \right\|_{L_2[0, 1]} + (L_1 + L_{05}) \left\| \frac{\partial^4 \varphi_3(x)}{\partial x^4} \right\|_{L_2[0, 1]} \right] |\omega_1 - \omega_2| + \\
& + M_0 \delta_0 \left(\frac{1}{\pi} \right)^2 \sqrt{\sum_{n=1}^{\infty} \frac{1}{n^4}} \left(|\nu| L_{04} \sqrt{\sum_{n=1}^{\infty} \frac{1}{|Z_n(\nu, \omega_1)|^2}} + \frac{|\nu|}{\sqrt[3]{\omega_2}} L_1 \sqrt{\sum_{n=1}^{\infty} \frac{1}{|Z_n(\nu, \omega_1)|^2}} \right) |\omega_1 - \omega_2| + \\
& \quad + \delta_0 M_0^2 \frac{|\nu|}{\sqrt[3]{\omega_2}} \left(\frac{1}{\pi} \right)^2 \sqrt{\sum_{n=1}^{\infty} \frac{1}{n^4}} \sqrt{\sum_{n=1}^{\infty} \frac{1}{|\bar{Z}_n(\nu)|^2}} \left[L_{04} + L_1 \frac{1}{\sqrt[3]{\omega_2}} \right] |\omega_1 - \omega_2| + \\
& + \alpha_0 \beta_0 \delta_2 M_0^2 \frac{|\nu|}{\sqrt[3]{\omega_2^2}} \left\| \frac{1}{\bar{\lambda}^2 \bar{Z}^2(\nu, \omega_2)} \right\|_{\ell_2} \left\| \int_0^T \int_0^1 |G(t, z)| \sum_{r=1}^{\infty} |a_r^T(t, \omega_1) - a_r^{T-1}(t, \omega_2)| b_r(z) dz dt \right\| + \\
& + M_0 T \left\| \frac{1}{\bar{\lambda}^2} \right\|_{\ell_2} \sqrt{\sum_{n=1}^{\infty} \left[\max_{0 \leq t \leq T} \left| \int_0^1 F(t, y, \cdot) b_n(y) dy \right| \right]^2} \left(L_{04} + L_1 \frac{1}{\sqrt[3]{\omega_2}} \right) |\omega_1 - \omega_2| + \\
& \quad + \frac{M_0}{\sqrt[3]{\omega_2}} \left\| \frac{1}{\bar{\lambda}^2} \right\|_{\ell_2} \sqrt{\sum_{n=1}^{\infty} \left| \int_0^1 l(y) b_n(y) dy \right|^2} \times \\
& \quad \times \int_0^T \int_0^1 |G(t, z)| \sum_{r=1}^{\infty} |a_r(t, \omega_1) - a_r(t, \omega_2)| b_r(z) dz dt, \quad (47)
\end{aligned}$$

where

$$\bar{Z}_n(\nu) = \frac{1}{\lambda_n} \begin{vmatrix} 1 - \nu \Delta_{11} & \nu \Delta_{12} & \dots & \nu \Delta_{1p} \\ \nu \Delta_{21} & 1 - \nu \Delta_{22} & \dots & \nu \Delta_{2p} \\ \dots & \dots & \dots & \dots \\ \nu \Delta_{p1} & \nu \Delta_{p2} & \dots & 1 - \nu \Delta_{pp} \end{vmatrix}, \quad \Delta_{ij} = \int_0^T \beta_i(s) \int_0^s \alpha_j(\theta) d\theta ds.$$

It is not difficult to check that from (47) we obtain

$$\|\bar{a}(t, \nu, \omega_1) - \bar{a}(t, \nu, \omega_2)\|_{B_2[0, T]} \leq M_4 |\omega_1 - \omega_2| + \rho \cdot \|\bar{a}(t, \nu, \omega_1) - \bar{a}(t, \nu, \omega_2)\|_{B_2[0, T]}, \quad (48)$$

where

$$\begin{aligned} M_4 = & C_0 M_0 \left(\frac{\sqrt{2}}{\pi} \right)^4 \sqrt{\sum_{n=1}^{\infty} \frac{1}{n^8}} \left[L_1 \left\| \frac{\partial^4 \varphi_1(x)}{\partial x^4} \right\|_{L_2[0,1]} + \right. \\ & \left. + (L_1 + L_{04}) \left\| \frac{\partial^4 \varphi_2(x)}{\partial x^4} \right\|_{L_2[0,1]} + (L_1 + L_{05}) \left\| \frac{\partial^4 \varphi_3(x)}{\partial x^4} \right\|_{L_2[0,1]} \right] + \\ & + |\nu| M_0 \delta_0 \left(\frac{1}{\pi} \right)^2 \sqrt{\sum_{n=1}^{\infty} \frac{1}{n^4}} \left[\sqrt{\sum_{n=1}^{\infty} \frac{1}{|Z_n(\nu, \omega_1)|^2}} + M_0 \frac{1}{\sqrt[3]{\omega_2}} \sqrt{\sum_{n=1}^{\infty} \frac{1}{|\bar{Z}_n(\nu)|^2}} \right] \left[L_{04} + \frac{1}{\sqrt[3]{\omega_2}} L_1 \right] + \\ & + T \left[M_0 L_{04} + \frac{L_1}{\sqrt[3]{\omega_2}} \right] \left\| \frac{1}{\lambda^2} \right\|_{\ell_2} \max_{0 \leq t \leq T} \|F(t, x, \cdot)\|_{L_2[0,1]}. \end{aligned}$$

From the estimate (48) we obtain (42). Theorem 4.2 is proved. \square

5. Convergence of the Fourier series

Theorem 5.1. *Let the conditions of the Theorem 3.1 be fulfilled. Then for regular values of the parameters ν the series (36) converges absolute and uniform in the domain Ω . Moreover, the solution of the mixed problem (1)–(3) belongs to the class of functions (4).*

The *proof* of the theorem 5.1 is based on obtaining the estimates for the Fourier series (36) and for its derivatives. The method for obtaining an estimate is the same as in the case of obtaining estimates (38)–(40), (47) and (48). The Theorem 5.1 is proved.

Corollary 5.2. *Let be fulfilled the conditions of the Theorem 3.1. Then the following estimate*

$$|V(t, \nu, \omega_1) - V(t, \nu, \omega_2)| \leq L_C |\omega_1 - \omega_2|, \quad 0 < L_C = \text{const}. \quad (49)$$

holds.

6. Conclusion

It is considered a fifth order nonlinear partial integro-differential equations (1) with mixed conditions (2) and (3) and with two real parameters ν, ω . The Fourier spectral method of separation of variables (5) is applied. A countable system of nonlinear functional-integral equations (35) is derived. Theorem on a uniqueness and existence of the solution of mixed problem (1)–(3) is proved for regular values of parameters. The method of compressing mapping is applied for countable system (35) in Banach space $B_2[0, T]$. The solution of the mixed problem (1)–(3) is obtained in the form of Fourier

series (36). Theorem on absolute and uniform convergence of Fourier series is proved. Continuous dependence on parameter ω of the classical solution of mixed problem is studied (see, estimates (42) and (49)).

We hope that this work can serve as a basis for further development of the theory of partial differential and integro-differential equations of the third and higher orders.

Support

This research was funded by the Science Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan (Grant No. AP23488740, 2024-2026).

References

- [1] V. M. Aleksandrov, and E. V. Kovalenko, *Problems of continuum mechanics with mixed boundary conditions*, Moscow, Nauka, 1986. 336 p. (in Russian)
- [2] G. Aman, *Nonlocal quasilinear parabolic equations*, Uspekhi Mat. Nauk, 2005, v. 60 (6 (366)), pp. 21–32.
- [3] S. N. Antontsev, A. V. Kazhikhov, and V. N. Monakhov, *Boundary value problems in the mechanics of inhomogeneous fluids*, Novosibirsk, SO Nauka, 1983. 319 p.
- [4] R. R. Ashurov, Yu. E. Fayziev, and N. Kh. Khushvaktov, *Forward and inverse problems for the Barenblatt–Zheltov–Kochina type fractional equations*, Lobachevskii J. Math., 2023, v. 44 (7), pp. 2563–2572.
- [5] R. R. Ashurov, and M. D. Shakarova, *Time-dependent source identification problem for fractional Schrodinger type equations*, Lobachevskii J. Math., 2022, v. 43 (5), pp. 1053–1064.
- [6] A. T. Assanova, *On solvability of nonlinear boundary value problems with integral condition for the system of hyperbolic equations*, Electronic J. Qualitative Theory of Differential Equat., 2015, v. 63, pp. 1–13.
- [7] A. T. Assanova, *One approach to the solution of a nonlocal problem for systems of hyperbolic equations with integral conditions*, J. Math. Sci. (N.Y.), 2019, v. 238 (3), pp. 189–206.
- [8] A. T. Assanova, *Solvability of a nonlocal problem for a hyperbolic equation with integral conditions*, Electronic J. Differential Equat., 2017, v. 2017 (170), pp. 1–12.
- [9] G. I. Barenblatt, Yu. P. Zheltov, and I. N. Kochina, *On finiteness conditions in the mechanics of continuous media. Static problems of the theory of elasticity*, Prikl. Mat. i Mekh., 1960, v. 24 (5), pp. 316–322. (in Russian)
- [10] G. I. Barenblatt, and Yu. P. Zheltov, *Fundamental equations of filtration of homogeneous liquids in fissured rocks*, Sov. Phys. Dokl., 1960, v. 132 (5), pp. 522–525.

- [11] G. I. Barenblatt, V. M. Yentov, and V. M. Ryzhik, *Movement of liquids and gases in natural reservoirs*, Moscow, Nedra, 1984 (in Russian).
- [12] K. S. Basniev, I. N. Kochina, and V. M. Maksimov, *Underground hydromechanics*, Moscow, Nedra, 1993 (in Russian).
- [13] K. Baykuziev, *Basic mixed problems for some degenerate partial differential equations*. Tashkent, Fan, 1984. 250 p. (in Russian)
- [14] D. J. Benney, and J. C. Luke, *Interactions of permanent waves of finite amplitude*, J. Math. Physics, 1964, **43**, pp. 309–313.
- [15] E. N. Bereslavskii, *Effect of evaporation or infiltration on the free surface of groundwater in certain problems of underground hydromechanics*, American J. Appl. Math. and Stat., 2017, v. 5 (5), pp. 159–163. <https://doi.org/10.12691/ajams-5-5-1>
- [16] A. N. Bogolyubov, and M. D. Malykh, *On one class of nonlocal nonlinear equations of parabolic type*, Zhurn. Vychisl. Matem. i Mat. Fiziki, 2011, v. 51 (6), pp. 1056–1063 (in Russian).
- [17] G. I. Chandirov, *Mixed problem for quasilinear equations of hyperbolic type*. Diss. ... Doc. Physics and Mathematics, Baku, Azerb. State Univ-ty, 1970. 248 p. (in Russian)
- [18] V. A. Chernyatin, *Justification of the Fourier method in a mixed problem for partial differential equations*, Moscow, MSU, 1992. 111 p. (in Russian)
- [19] M. M. Cavalcanti, V. N. Domingos Cavalcanti, and J. Ferreira, *Existence and uniform decay for a nonlinear viscoelastic equation with strong damping*, Math. Methods in the Applied Sciences, 2001, v. 24, pp. 1043–1053.
- [20] Yu. A. Dubinsky, *Quasilinear elliptic and parabolic equations of some order*, Uspekhi Mat. Nauk, 1968, v. 23 (1), pp. 45–90 (in Russian).
- [21] D. G. Gordesiani, G. A. Avalishvili, *Solutions of nonlocal problems for one-dimensional oscillations of a medium*, Mat. Modelirovanie, 2000, v. 12 (1), pp. 94–103 (in Russian).
- [22] O. M. Guseynov, and T. J. Mamedov, *Fundamental solution of a Cauchy and Goursat type problem for second order hyperbolic equations and related optimal control problems*, Journal of Contemporary Applied Mathematics, 2024, v. 14 (1), pp. 70–77. <https://doi.org/10.5281/zenodo.10908206>
- [23] V. B. Dmitriev, *Nonlocal problem with integral conditions for the wave equation*, Vestnik SamGU. Estestvennonauchnaya seriya, 2006, v. 42 (2), pp. 15–27 (in Russian).
- [24] V. A. Ilyin, *On the solvability of mixed problems for hyperbolic and parabolic equations*, Uspekhi Mat. Nauk, 1960, v. 15 (2), pp. 97–154 (in Russian).

- [25] N. I. Ivanchov, *Boundary value problems for a parabolic equation with an integral condition*, Differents. Uravneniya, 2004, v. 40 (4), pp. 547–564 (in Russian).
- [26] N. I. Ionkin, *Solution of a boundary value problem in the theory of heat conduction with a non-classical boundary condition*, Differents. Uravneniya, 1977, v. 13 (2), pp. 294–304 (in Russian).
- [27] I. E. Egorov, and E. S. Efimova, *Modified Galerkin method for a semilinear parabolic equation with changing direction of time*, Sibir. Zhurn. Chistoy i Prikladnoy Mat., 2016, v. 16 (2), pp. 6–15 (in Russian).
- [28] M. O. Korpusov, *Destruction in parabolic and pseudoparabolic equations with double nonlinearities*, Moscow, URSS, 2012. 184 p. (in Russian)
- [29] O. A. Ladyzhenskaya, V. A. Solonnikov, and N. N. Uraltseva, *Linear and quasilinear equations of parabolic type*, Moscow, Nauka, 1967. 756 p. (in Russian)
- [30] V. P. Mikhailov, *Partial differential equations*, Moscow, Nauka, 1976. 391 p. (in Russian)
- [31] A. M. Nakhushev, *Nonlocal problem and Goursat problem for a loaded equation of hyperbolic type and their applications to the forecast of soil moisture*, Doklady AN SSSR, 1978, v. 242 (5), pp. 1008–1011. (in Russian)
- [32] P. Ya. Polubarinova-Kochina, *The theory of groundwater movement*, Moscow, Nauka, 1977 (in Russian).
- [33] A. D. Polyanin, V. F. Zaitsev, and A. I. Zhurov, *Methods for solving nonlinear equations of mathematical physics and mechanics*, Moscow, Fizmatlit, 2005. 256 p. (in Russian)
- [34] L. S. Pulkina, *Mixed problem with an integral condition for a hyperbolic equation*, Mat. Zametki, 2003, v. 74 (3), pp. 435–445.
- [35] M. A. Sagadeeva, and F. L. Hasan, *Bounded solutions of Barenblatt–Zhel'tov–Kochina model in Quasi-Sobolev spaces*, Bulletin of the South Ural State University. Ser. Math. Modelling, Programming and Computer Software, 2015, v. 8 (4), pp. 138–144.
- [36] K. Kh. Shabadikov, *Study of solutions of mixed problems for quasilinear differential equations with a small parameter on the highest mixed derivative*. Diss. . . . Cand. Physics and Mathematics. Fergana State Univ-ty, Fergana, 1984. 156 p. (in Russian)
- [37] V. A. Steklov, *Basic problems of mathematical physics*. Moscow, Nauka, 1983. 433 p. (in Russian)
- [38] G. A. Sviridyuk, and D. E. Shafranov, *The Cauchy problem for the Barenblatt–Zhel'tov–Kochina equation on a smooth manifold*, Vest. ChelyabGU, 2003, no 9, pp. 171–177.

- [39] Zh. O. Takhirov, and R. N. Turaev, *On one nonlocal problem for a nonlinear parabolic equation*, Vladikavkaz. Mat. Zhurn., 2014, v. 16 (1), pp. 42–49 (in Russian).
- [40] A. I. Vagabov, *Generalized Fourier method for solving mixed problems for nonlinear equations*, Differ. Equat., 1996, v. 32 (1), pp. 94–105.
- [41] T. K. Yuldashev, *Mixed value problem for a nonlinear differential equation of fourth order with small parameter on the parabolic operator*, Comput. Math. Math. Phys., 2011, v. 51 (9), pp. 1596–1604.
- [42] T. K. Yuldashev, *Mixed value problem for nonlinear integro-differential equation with parabolic operator of higher power*, Comput. Math. Math. Phys., 2012, v. 52 (1), pp. 105–116.
- [43] T. K. Yuldashev, *Nonlinear optimal control of thermal processes in a nonlinear inverse problem*, Lobachevskii J. Math., 2020, v. 41 (1), pp. 124–136.
- [44] T. K. Yuldashev, *Solvability of a boundary value problem for a differential equation of the Boussinesq type*, Differential Equat., 2018, v. 54 (10), pp. 1384–1393.
- [45] T. K. Yuldashev, and A. K. Fayziyev, *Mixed problem for a linear differential equation of parabolic type with nonlinear impulsive conditions*, Nanosystems: Physics. Chemistry. Mathematics, 2024, v. 15 (2), pp. 160–169. <https://doi.org/10.17586/2220-8054-2024-15-2-160-169>
- [46] T. K. Yuldashev, A. K. Fayziyev, and F. D. Rakhmonov, *Mixed problem for a nonlinear impulsive differential equation of parabolic type*, Chelyab. Phys.-Math. J., 2024, v. 9, (1), pp. 111–123.
- [47] T. K. Yuldashev, B. I. Islomov and E. K. Alikulov, *Boundary value problems for loaded third-order parabolic-hyperbolic equations in infinite three-dimensional domains*, Lobachevskii J. Math., 2020, v. 41 (5), pp. 926–944.
- [48] T. K. Yuldashev, and B. J. Kadirkulov, *Inverse problem for a partial differential equation with Gerasimov–Caputo-type operator and degeneration*, Fractal Fract., 2021, v. 5 (2), ID 58, pp. 1–13. <https://doi.org/10.3390/fractalfract5020058>
- [49] T. K. Yuldashev and K. Kh. Shabadikov, *Mixed problem for a higher-order nonlinear pseudoparabolic equation*, J. Math. Sci. (N.Y.), 2021, v. 254 (6), pp. 776–787.
- [50] T. K. Yuldashev and K. Kh. Shabadikov, *Initial value problem for a higher order quasilinear partial differential equation*, J. Math. Sci. (N.Y.), 2021, v. 254 (6), pp. 811–822.
- [51] N. P. Vekua, *Some Questions in the theory of differential equations and applications in mechanics*. Moscow, Nauka, 1991. 256 p.

- [52] A. Anguraj, A. Vinodkumar, *Global existence and stability results for partial delay integro-differential equations with random impulses*, *Filomat*, 2023, v. 37 (1), pp. 317–334. <https://doi.org/10.2298/FIL2301317A>
- [53] Zh. A. Artykova, R. A. Bandaliyev, and T. K. Yuldashev, *Nonlocal direct and inverse problems for a second order nonhomogeneous Fredholm integro-differential equation with two redefinition data*, *Lobachevskii J. Math.*, 2023, v. 44 (10), pp. 4215–4230. <https://doi.org/10.1134/S1995080223100050>
- [54] S. N. Askhabov, *On a second-order integro-differential equation with difference kernels and power nonlinearity*, *Bulletin of the Karaganda University. Mathematics series*, 2022, v. 106 (2), pp. 38–48. DOI 10.31489/2022M2/38-48
- [55] A. T. Assanova, *A two-point boundary value problem for a fourth order partial integro-differential equation*, *Lobachevskii J. Math.*, 2021, v. 42 (3), pp. 526–535. <https://doi.org/10.1134/S1995080221030082>
- [56] A. T. Assanova and S. N. Nurmukanbet, *A solvability of a problem for a Fredholm integro-differential equation with weakly singular kernel*, *Lobachevskii J. Math.*, 2022, v. 43 (1), pp. 182–191. <https://doi.org/10.1134/S1995080222040047>
- [57] N. Aviltay and M. Akhmet, *Asymptotic behavior of the solution of the integral boundary value problem for singularly perturbed integro-differential equations*, *J. Mathematics, Mechanics and Computer Science*, 2021, v. 112 (4), pp. 13–28. <https://doi.org/10.26577/JMMCS.2021.v112.i4.02>
- [58] A. Bellour and Kh. Rouiban, *Iterative continuous collocation method for solving non-linear Volterra integro-differential equations*, *Proceedings of the Institute of Mathematics and Mechanics. National Academy of Sciences of Azerbaijan*, 2021, v. 47 (1), pp. 99–111. <https://doi.org/10.30546/2409-4994.47.1.99>
- [59] V. F. Chistyakov and E. V. Chistyakova, *Properties of degenerate systems of linear integro-differential equations and initial value problems for these equations*, *Differential Equations*, 2023, v. 59 (1), pp. 13–28. <https://doi.org/10.1134/S0012266123010023>
- [60] D. K. Durdiev and J. S. Safarov, *Finding the two-dimensional relaxation kernel of an integro-differential wave equation*, *Differential Equations*, 2023, v. 59 (2), pp. 214–229. <https://doi.org/10.1134/S0012266123020064>
- [61] S. Iskandarov, *Lower bounds for the solutions of a first-order linear homogeneous Volterra integro-differential equation*, *Differential Equations*, 1995, v. 31 (9), pp. 1462–1466.
- [62] S. Iskandarov and G. T. Khalilov, *On lower estimates of solutions and their derivatives to a fourth-order linear integrodifferential Volterra equation*, *J. Math. Sci. (N. Y.)*, 2018, v. 230 (5), pp. 688–694.

- [63] I. G. Mamedov, and A. J. Abdullayeva, *On correct solvability of the boundary value problem in the non-classical treatment set on the middle of the domain for one integrodifferential 3D Bianchi equation*, Journal of Contemporary Applied Mathematics, 2018, v. 8 (1), pp. 69–80. (in Russian)
- [64] I. G. Mamedov, and A. J. Abdullayeva, *Fundamental solution of one boundary value problem on the geometric middle of the domain for the 3d Bianchi integro-differential equation*, Journal of Contemporary Applied Mathematics, 2024, v. 14 (2), pp. 26–37. <https://doi.org/10.69624/2222-5498.2024.2.26>
- [65] N. A. Rautian and V. V. Vlasov, *Spectral analysis of the generators for semigroups associated with Volterra integro-differential equations*, Lobachevskii J. Math., 2023, v. 44 (3), pp. 926–935.
- [66] Yu. G. Smirnov, *On the equivalence of the electromagnetic problem of diffraction by an inhomogeneous bounded dielectric body to a volume singular integro-differential equation*, Comput. Math. and Math. Physics, 2016, v. 56 (9), pp. 1631–1640.
- [67] V. B. Vasilyev and Sh. H. Kutaiba, *Asymptotic behavior of solution to some boundary value problem*, Azerbaijan J. Math., 2023, v. 13 (1), pp. 51–61.
- [68] S. Yaghoubi, H. Aminikhaha, and Kh. Sadri, *An effective operational matrix method based on shifted sixth-kind Chebyshev polynomials for solving fractional integro-differential equations with a weakly singular kernel*, Filomat, 2024, v. 38 (7), pp. 2457–2486. <https://doi.org/10.2298/FIL2407457Y>
- [69] A. Yakar and H. Kutlay, *Extensions of some differential inequalities for fractional integro-differential equations via upper and lower solutions*, Bulletin of the Karaganda University. Mathematics series, 2023, v. 109 (1), pp. 156–167. <https://doi.org/10.31489/2023M1/156-167>
- [70] V. A. Yurko, *Inverse problems for first-order integro-differential operators*, Math. Notes, 2016, v. 100 (6), pp. 876–882.
- [71] T. K. Yuldashev, *Nonlocal inverse problem for a pseudohyperbolic-pseudoelliptic type integro-differential equations*, Axioms, 2020, v. 9 (2), ID 45, pp. 1–21. <https://doi.org/10.3390/axioms9020045>
- [72] T. K. Yuldashev, *On a boundary value problem for a fifth order partial integro-differential equation*, Azerbaijan J. Math., 2022, v. 12 (2), pp. 154–172.
- [73] T. K. Yuldashev, Z. K. Eshkuvatov and N. M. A. Nik Long, *Nonlinear Fredholm functional-integral equation of first kind with degenerate kernel and integral maxima*, Malaysian Journal of Fundamental and Applied Sciences, 2023, v. 19, pp. 82–92.

- [74] T. K. Yuldashev and A. K. Fayziyev, *Inverse problem for a second order impulsive system of integro-differential equations with two redefinition vectors and mixed maxima*, *Nanosystems: Physics. Chemistry. Mathematics*, 2023, v. 14 (1), pp. 13–21. <https://doi.org/10.17586/2220-8054-2023-14-1-13-21>
- [75] T. K. Yuldashev, and O. Sh. Kilichev, *Inverse problem for a hyperbolic integro-differential equation with two redefinition conditions at the end of the interval and involution*, *Azerb. J. Math.*, 2024, v. 14 (1), pp. 3–22. <https://doi.org/10.59849/2218-6816.2024.1.3>
- [76] T. K. Yuldashev, F. D. Rakhmonov, *On a Benney–Luke type differential equation with nonlinear boundary value conditions*, *Lobachevskii J. Math.*, 2021, v. 42 (15), pp. 3761–3772. <https://doi.org/10.1134/S1995080222030210>
- [77] G. V. Zavizion, *Asymptotic solutions of systems of linear degenerate integro-differential equations*, *Ukr. Math. J.*, 2003, v. 55 (4), pp. 521–534.
- [78] T. K. Yuldashev and S. K. Zarifzoda, *On a new class of singular integro-differential equations*, *Bulletin of the Karaganda University. Mathematics series*, 2021, v. 101 (1), pp. 138–148. <https://doi.org/10.31489/2021M1/138-148>
- [79] A. A. Boichuk and A. P. Strakh, *Noetherian boundary-value problems for systems of linear integro-dynamical equations with degenerate kernel on a time scale*, *Nelineynye kolebaniya*, 2014, v. 17 (1), pp. 32–38. (in Russian)
- [80] D. S. Djumabaev and E. A. Bakirova, *On one single solvability of boundary value problem for a system of Fredholm integro-differential equations with degenerate kernel*, *Nelineynye kolebaniya*, 2015, v. 18 (4), pp. 489–506. (in Russian)
- [81] T. K. Yuldashev, *Mixed problem for pseudoparabolic integro-differential equation with degenerate kernel*, *Differential Equations*, 2017, v. 53 (1), pp. 99–108.
- [82] T. K. Yuldashev, *Nonlocal mixed-value problem for a Boussinesq-type integro-differential equation with degenerate kernel*, *Ukrain. Math. J.*, 2017, v. 68 (8), pp. 1278–1296.
- [83] T. K. Yuldashev, *On a boundary value problem for a fifth order partial integro-differential equation*, *Azerb. J. Math.*, 2022, v. 12 (2), pp. 154–172.
- [84] T. K. Yuldashev, *Inverse problem for a nonlinear Benney–Luke type integro-differential equations with degenerate kernel*, *Russian Math. (Izv. vuz)*, 2016, v. 60 (8), pp. 53–60.
- [85] T. K. Yuldashev, *Inverse boundary-value problem for an integro-differential Boussinesq-type equation with degenerate kernel*, *J. Math. Sci. (NY)*, 2020, v. 250 (5), pp. 847–858. <https://doi.org/10.1007/s10958-020-05050-2>

- [86] T. K. Yuldashev, *Determining of coefficients and the classical solvability of a non-local boundary-value problem for the Benney–Luke integro-differential equation with degenerate kernel*, J. Math. Sci. (NY), 2021, v. 254 (6), pp. 793–807.
- [87] T. K. Yuldashev, Yu. P. Apakov and A. Kh. Zhuraev, *Boundary value problem for third order partial integro-differential equation with a degenerate kernel*, Lobachevskii J. Math., 2021, v. 42 (6), pp. 1317–1327.
- [88] T. K. Yuldashev, Zh. A. Artykova, and Sh. U. Alladustov, *Nonlocal problem for a second order Fredholm integro-differential equation with degenerate kernel and real parameters*, Proceedings of the Institute of Mathematics and Mechanics of National Academy of Sciences of Azerbaijan, 2023, v. 49 (2), pp. 228–242. <https://doi.org/10.30546/2409-4994.2023.49.2.228>

Tursun K. Yuldashev
Tashkent State Transport University, Alley Temiryulchilar 1, Tashkent, Uzbekistan
E-mail: tursun.k.yuldashev@gmail.com

Khanlar R. Mamedov
İğdir university, İğdir, Turkey
E-mail: hanlar.residoglu@igdir.edu.tr

Mahkambek M. Babayev
Tashkent State Transport University, Alley Temiryulchilar 1, Tashkent, Uzbekistan
E-mail: babayevm@mail.ru

Received 10 June 2024

Accepted 17 October 2024